

AstaREAL® Astaxanthin

New AstaREAL® Astaxanthin Study Shows Improved Lipid Metabolism or “Fat-Burn” combined with Exercise

After the Christmas and New Year Festivities, people tend to gain a few pounds hoping to lose it after the holiday season. Resolutions may include a promise to the gym but as we all know too well, good intentions and will-power quickly diminish. A new study published by Japanese scientists suggests that help may be at hand to support that resolution.

In the past, **AstaREAL®** astaxanthin research had demonstrated improved muscle endurance†, reduced lactic acid build-up, lowered membrane peroxidation and many more. Currently, supplements containing **AstaREAL®** are used by several world-class athletes from around the globe. New findings by Aoi *et al.*, (2008) reveal an opening into the mechanism of action behind increased endurance and at the same time show new indications of enhanced fat metabolism in mice model.

The most recent data published by the researchers from the Kyoto Prefectural University, University of Shizuoka, University of Hyogo, and University of Nagoya revealed that astaxanthin protected the carnitine palmitoyl transferase I (CPT I) function in mitochondria in mice during intense physical activity (treadmill 30 m/minute). Furthermore, the astaxanthin treated group accelerated body fat reduction or “fat-burning” when combined with exercise compared to just exercise alone. The



authors suggested enhanced lipid metabolism during physical activity may be the reason behind increased endurance and body fat reduction†.

The CPT I is a lipid transport enzyme is located on the mitochondrial membrane and it supplies lipids or “fuel” into mitochondria for energy production. During intense physical activity muscle mitochondria produce high amounts of reactive oxygen species (ROS) that could lead to CPT I oxidation resulting in decreased function of lipid transport. The results showed that **AstaREAL®** astaxanthin protected CPT I against ROS which subsequently allowed the continual transport of fats into the mitochondria for energy production ($p < 0.05$). Evidence include decreased CPT I modification by HEL lipid peroxide in Figure 1. ($p < 0.05$) and significantly increased colocalization of Fat/CD36 with CPT I ($p < 0.05$) on the mitochondrial membrane (not shown).

Fat utilization compared to carbohydrates was the predominant source of ATP energy according to the Respiratory Exchange Ratio (RER). Furthermore, plasma lactate significantly increased by exercise, while this elevation was suppressed by **AstaREAL®** astaxanthin in the diet ($P < 0.05$). As a result, the time to reach exhaustion significantly increased by 20% longer than the control group ($p < 0.05$); refer to Figure 2. After 4 weeks of **AstaREAL®** intake, the body weight and epididymal fat showed a significant decrease compared to the exercise control and sedentary groups (Table 1).




Figure 2. Running time to exhaustion. 30m/min after 4 weeks of **AstaREAL®** astaxanthin intake. Mean values, N=8, ($p < 0.05$) compared to control diet.




Figure 1. Analysis of CPT I modification by HEL lipid peroxide. Lysate protein from the muscle collected after running. Immunoprecipitated with CPT I antibody, SDS-PAGE separated and probed for HEL. Mean values, N=6

	Sedentary		Training	
	Control	AstaREAL®	Control	AstaREAL®
Body weight (g)	48.0 ± 1.1	46.8 ± 1.1	42.5 ± 1.0*	40.0 ± 1.0*
Gastrocnemius muscle (mg/g BW)	9.9 ± 0.3	9.7 ± 0.3	9.7 ± 0.2	10.0 ± 0.2
Epididymal fat (mg/g BW)	51.4 ± 2.2	48.3 ± 3.0	43.5 ± 2.0*	36.6 ± 2.0*#

Table 1. Body weight and tissue weight. Effects of training and AstaREAL® treatment on body composition. Mean values, N=8. *Compared to sedentary (p<0.05); # compared to control diet (p<0.05).

This study is important because it showed that **AstaREAL®** astaxanthin can modify muscle metabolism by its antioxidant property, resulting in improved muscle performance and weight loss benefits. Recent studies using **AstaREAL®** in endurance studies also point to the same findings by Aoi *et al.*, 2008. In September 2007 at the 21st Annual Meeting on Carotenoid Research, Osaka, Japan; Professor Yazawa of Tokyo University of Marine Science and Technology presented similar findings of improved lipid metabolism with exercise on a diet with **AstaREAL®** astaxanthin and high fat intake. Yazawa's research group has also published other papers on this subject. The relevant publications are Ikeuchi *et al.*, 2006 Effects of astaxanthin supplementation on exercise-induced fatigue in mice and Ikeuchi *et al.*, 2007 Effects of astaxanthin in obese mice fed a high-fat diet.

Reference: 2008 Aoi *et al.*, Astaxanthin improves muscle lipid metabolism in exercise via inhibitory effect of oxidative CPT I modification. *Biochemical & Biophysical Research Communications* 36:892-897.

†Pat US PAT 6245818 and other countries

‡Pat Pending

Fuji Chemical Industry, Japan is the world leader in the production and research of **AstaREAL®** natural astaxanthin. For more information, please visit www.astareal.com or [contact us](#).

AstaREAL is a trademark or registered trademark of Fuji Chemical Industry Co., Ltd. in Japan, United States of America, Europe and/or other countries.

FUJI CHEMICAL INDUSTRY CO., LTD.

For more information, please [contact us](#). If you would prefer not to receive email newsletter, or you've changed your email address, please [click here](#).

Information found is presented in good faith and no guarantee or obligation as to accuracy and no assumption of liability. The information herein does not imply a performance warranty. Purchasers should decide the suitability of the product with respect to their desired application and purpose.